



COLONOSCOPY PREPARATION

*For the best exam possible, your colon must be clean. Your preparation will begin the day before your procedure.
Please read the instructions carefully before you begin.*

-PENDING CLEARANCE: Hold blood thinners (Plavix, Xarelto, Eliquis, Effient, Coumadin/Warfarin) **THREE TO FIVE DAYS** before the procedure(s).

-Hold Aspirin 325 mg **FIVE DAYS** before the procedure. -Hold iron **FIVE DAYS** before the procedure.

-Hold Vitamins C and E **TWO DAYS** before the procedure.

-Hold anti-inflammatory medication **TWO DAYS** before the procedure. (Tylenol is okay to take.)

If **DIABETIC**, please take half a dose of your Diabetic medication starting from the day before your procedure(s).
If further questions, please refer to your Primary Care Provider

ATTENTION: Please **STOP** following medications **7 days** before procedures:

-OZEMPIC

- BYETTA

-ADLYXIN

-BYDUREON

-RYBELSUS

-TRULICITY

-VICTOZA

-SAXENDA

-WEGOVY

-ZEPBOUND

-MOUNJARO

**STARTING ON _____ / _____ / _____ ALL DAY (THE DAY
BEFORE PROCEDURE(S))**

CLEAR LIQUID DIET

CLEAR LIQUIDS INCLUDE:

- Water
- Broth/Bouillon
- Coconut water (**NO PULP**)
- Clear Juices (**NO PULP**) - Apple, **WHITE** Cranberry, **WHITE** Grape
- Coffee (**BLACK** - **NO** creamer, **NO** dairy products)
- Tea (**HOT** or **COLD**) sugar or honey is okay (**NO** dairy products)
- Sports drinks - clear only, **NO** green, purple or red
- Clear carbonated soft drinks - Sprite, 7UP and/or Ginger Ale
- Yellow gelatin (**NO** toppings) - **STOP AFTER 5:00 P.M. (DAY BEFORE THE PROCEDURE)**
- You may use menthol lozenges to help tolerate preparation (Halls cough drops - **CLEAR ONLY**)
- ★ **DO NOT have dark liquids (red, orange, green, blue or purple)**
- ★ **DO NOT consume solid foods, dairy products, and cloudy liquids**
- ★ **STOP all clear liquids after midnight except water**

*If you have a dry mouth, you may rinse with water and spit out. **DO NOT** swallow the water. If your stomach is not empty, you may feel nauseated with the sedation and may vomit causing pneumonia.

100 S. ANAHEIM BLVD SUITE #250 ANAHEIM, CA 92805
955 W. IMPERIAL HWY SUITE # 100 BREA, CA 92821
TELE (714) 527-6000 FAX (714) 527-2371

BOWEL PREPARATIONS INSTRUCTIONS

GENERIC (1 GALLON) BOWEL PREP FOR
COLONOSCOPY SPLIT DOSE

- You may have as much clear liquids as desired until 12 midnight, **NOTHING BY MOUTH AFTER MIDNIGHT EXCEPT THE SECOND DOSAGE OF BOWEL PREP.** An empty stomach is necessary for a successful COLONOSCOPY

/ /202_	/ /202_
STEP ONE DAY BEFORE PROCEDURE <u>5 P.M.</u> start your bowel prep. <u>Drink 8 oz.</u> <u>glass every 15 minutes</u> for a total of 12 - <u>8 oz. glasses</u>  5:00  6:00  7:00  5:15  6:15  7:15  5:30  6:30  7:30  5:45  6:45  7:45	STEP TWO 5 HOURS BEFORE YOUR PROCEDURE ____: ____ <u>A.M.</u> Drink remaining prep. <u>Drink 8 oz.</u> <u>glass every 15 minutes</u> for a total of 4 - <u>8 oz. glasses</u>  :  :  :  :
<p><u>PLEASE PICK UP MEDICATION AT YOUR PHARMACY WITHIN 7 DAYS</u> If you have any questions regarding the bowel prep instructions, please contact our office at: (714) 527-6000</p>	

!!!PLEASE DO NOT THROW AWAY THE LEFTOVERS!!!